

# BencoCaine™ Singles

Topical Anesthetic Gel

## INSTRUCTIONS FOR USE

59° – 86° F  
15° – 30° C



**CHERRY** [4902-465]

Contains 120 Singles (0.3ml unit doses)

**NDC 60640-9000**

### DRUG FACTS

#### Active ingredient:

Benzocaine 20%

#### Purpose:

Oral Anesthetic

#### Uses:

BencoCaine is a flavored topical anesthetic gel featuring rapid onset, no systemic absorption and a 15 minute duration. BencoCaine is packaged in single-dose packages for maximum asepsis and ease of use. BencoCaine is also ideal for take-home, post-operative topical pain relief. Each package contains a single dose of BencoCaine plus swab applicator.

For the temporary relief of occasional minor irritation and pain, associated with:

- Sore mouth and throat
- Ulcers
- Minor dental procedures
- Needle punctures
- Minor irritation of the mouth and gums caused by dentures or orthodontic appliances

#### Warnings:

For external use only

#### Methemoglobinemia Warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- Pale, gray or blue colored skin (cyanosis)
- Headache
- Rapid heart rate
- Shortness of breath
- Dizziness or lightheadedness
- Fatigue or lack of energy

#### Allergy Alert:

Do not use if you have a history of allergy to any “caine” anesthetics, such as benzocaine, butacaine, procaine or tetracaine.

#### Sore Throat Warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, see your dentist or doctor promptly.

#### Do Not Use:

- For teething
- In children under 2 years of age

When using this product avoid contact with eyes

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center immediately.

#### Directions:

Remove one BencoCaine package from the 8-unit tray. Slowly peel back the lid, exposing the cotton swab applicator and unit dose well of BencoCaine. Dip swab applicator and apply gel to mucosa or gingival soft tissue. Wait one minute or until tissue feels numb before starting needle injection or other procedure. After treatment, one or more BencoCaine applications may be used for post-operative pain or discomfort.

**Do not exceed recommended dosage.**

#### Adults and children 2 years of age and older:

Apply to affected area up to 4 times daily or as directed by dentist or doctor.

#### Children under 12 years of age:

Should be supervised in the use of the product.

**Children under 2 years of age:** Do not use

#### Other information:

Store at 59° to 86° F (15° to 30° C)

#### Inactive ingredients:

Blue 1\*, flavor, PEG-8, PEG-75, Red 3\*, Red 33\*, Red 40\*, sodium saccharin, Yellow 10\*

\*Contains one or more of these ingredients.

#### Questions or Comments?

Distributed by

**BencoDental™**

800.GO.BENCO • [benco.com](http://benco.com)

295 CenterPoint Blvd., Pittston, PA 18640